# Doxorubicin

Section: 8. Immunomodulators and antineoplastics > 8.2. Antineoplastics and supportive medicines > 8.2.1. Cytotoxic medicines

**EMLc** Codes ATC: L01DB01

| | |
|---|---|
| **Indication** | Malignant neoplasms of kidney, except renal pelvis Code ICD11: 2C90.Z |
| **INN** | Doxorubicin |
| **Type de médicament** | Chemical agent |
| **Type de liste** | Liste complémentaire |
| **Formulations** | **Parenteral > General injections > IV:** 10 mg in vial (hydrochloride) powder for injection ; 50 mg in vial (hydrochloride) powder for injection ; 2 mg per mL in 5 mL vial (hydrochloride) ; 2 mg per mL in 25 mL vial (hydrochloride) |
| **Historique des statuts LME** | Ajouté pour la première fois en 2015 (TRS 994)<br>Modifié en 2023 (TRS 1049) |
| **Sexe** | Tous |
| **Âge** | Aussi recommandé pour les enfants |
| **Équivalence thérapeutique** | La recommandation concerne ce médicament spécifique |
| **Renseignements sur le brevet** | Patents have expired in most jurisdictions<br>Lire la suite sur les brevets. |
| **Balises** | Cancer |
| **Wikipédia** | Doxorubicin |
| **DrugBank** | Doxorubicin |

## Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended the inclusion of an additional dose forms of doxorubicin (2 mg/mL (hydrochloride) in 5 mL, 25 mL vial) to the EML and EMLc.