Doxorubicin

Section: 8. Immunomodulators and antineoplastics > 8.2. Antineoplastics and supportive medicines > 8.2.1. Cytotoxic medicines

EMLc ATC codes: L01DB01

Indication	Burkitt lymphoma including Burkitt leukaemia ICD11 code: 2A85.6
INN	Doxorubicin
Medicine type	Chemical agent
List type	Complementary
Formulations	**Parenteral > General injections > IV:** 10 mg in vial (hydrochloride) powder for injection ; 50 mg in vial (hydrochloride) powder for injection ; 2 mg per mL in 5 mL vial (hydrochloride) ; 2 mg per mL in 25 mL vial (hydrochloride)
EML status history	First added in 2011 (TRS 965) Changed in 2015 (TRS 994) Changed in 2023 (TRS 1049)
Sex	All
Age	Also recommended for children
Therapeutic alternatives	The recommendation is for this specific medicine
Patent information	Patents have expired in most jurisdictions Read more about patents.
Tags	Cancer
Wikipedia	Doxorubicin
DrugBank	Doxorubicin

Summary of evidence and Expert Committee recommendations

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended the inclusion of an additional dose forms of doxorubicin (2 mg/mL (hydrochloride) in 5 mL, 25 mL vial) to the EML and EMLc.