Doxorubicin

Section: 8. Immunomodulators and antineoplastics > 8.2. Antineoplastics and supportive medicines > 8.2.1. Cytotoxic medicines

EMLc | Codes ATC: L01DB01

Indication	Burkitt lymphoma including Burkitt leukaemia — Code ICD11: 2A85.6
INN	Doxorubicin
Type de médicament	Chemical agent
Type de liste	Liste complémentaire
Formulations	**Parenteral > General injections > IV:** 10 mg in vial (hydrochloride) powder for injection ; 50 mg in vial (hydrochloride) powder for injection ; 2 mg per mL in 5 mL vial (hydrochloride) ; 2 mg per mL in 25 mL vial (hydrochloride)
Historique des statuts LME	Ajouté pour la première fois en 2011 (TRS 965) Modifié en 2015 (TRS 994) Modifié en 2023 (TRS 1049)
Sexe	Tous
Âge	Aussi recommandé pour les enfants
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets.
Balises	Cancer
Wikipédia	Doxorubicin
DrugBank	Doxorubicin

Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended the inclusion of an additional dose forms of doxorubicin (2 mg/mL (hydrochloride) in 5 mL, 25 mL vial) to the EML and EMLc.