# Cyclophosphamide

Section: 8. Immunomodulators and antineoplastics > 8.2. Antineoplastics and supportive medicines > 8.2.1. Cytotoxic medicines

**EMLc** Codes ATC: L01AA01

| | |
|---|---|
| **Indication** | Rhabdomyosarcoma primary site Code ICD11: 2B55.Z |
| **INN** | Cyclophosphamide |
| **Type de médicament** | Chemical agent |
| **Type de liste** | Liste complémentaire |
| **Formulations** | **Parenteral > General injections > IV:** 500 mg in vial powder for injection ; 1 g in vial powder for injection ; 2 g in vial powder for injection<br>**Oral > Solid > dosage form:** 25 mg ; 50 mg |
| **Historique des statuts LME** | Ajouté pour la première fois en 1977 (TRS 615)<br>Modifié en 2015 (TRS 994)<br>Modifié en 2019 (TRS 1021)<br>Modifié en 2021 (TRS 1035)<br>Modifié en 2023 (TRS 1049) |
| **Sexe** | Tous |
| **Âge** | Aussi recommandé pour les enfants |
| **Équivalence thérapeutique** | La recommandation concerne ce médicament spécifique |
| **Renseignements sur le brevet** | Patents have expired in most jurisdictions<br>Lire la suite sur les brevets. |
| **Balises** | Cancer |
| **Wikipédia** | Cyclophosphamide |
| **DrugBank** | Cyclophosphamide |

## Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended the listings for oral dosage forms of cyclophosphamide on the EML and EMLc be amended to replace 'tablet' with 'solid oral dosage form' to reflect the availability also of capsules and allow greater flexibility in procurement.