Rifampicin

Section: 6. Anti-infective medicines > 6.2. Antibacterials > 6.2.4. Antileprosy medicines

EMLc | Codes ATC: J04AB02

Indication	Leprosy Code ICD11: 1B20.Z
INN	Rifampicin
Type de médicament	Chemical agent
Groupes d'antibiotiques	WATCH
Type de liste	Liste de base
Additional notes	Medicines for leprosy should be used in accordance with recommendations in WHO guidelines. Medicines used in the treatment of leprosy should never be used except in combination. Combination therapy is essential to prevent the emergence of drug resistance. Colour-coded blister packs (MDT blister packs) containing standard two-medicine (paucibacillary leprosy) or three-medicine (multibacillary leprosy) combination for adult and childhood leprosy should be used. MDT blister packs can be supplied free of charge through WHO.
Formulations	**Oral > Liquid:** 20 mg per mL (EMLc) **Oral > Solid:** 150 mg ; 300 mg
Historique des statuts LME	Ajouté pour la première fois en 1977 (TRS 615) Modifié en 1982 (TRS 685) Modifié en 2003 (TRS 920) Modifié en 2007 (TRS 950) Modifié en 2023 (TRS 1049) Modifié en 2025 (TRS 1064)
Sexe	Tous
Âge	Aussi recommandé pour les enfants
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets.
Wikipédia	Rifampicin
DrugBank	Rifampicin

Résumé des preuves et recommandation du comité d'experts

Addition of a note stating medicines for leprosy should be used in accordance with recommendations in WHO guidelines.