

EMLc

Codes ATC: J01CA04

<b>Indication</b>	Acute pharyngitis <span style="border: 1px solid #ccc; padding: 2px;">Code ICD11: CA02</span>
<b>INN</b>	Amoxicillin
<b>Type de médicament</b>	Chemical agent
<b>Groupes d'antibiotiques</b>	<span style="border: 1px solid #ccc; border-radius: 50%; padding: 2px 5px; color: #fff; background-color: #007bff;">A</span> ACCESS
<b>Type de liste</b>	Liste de base (EML) (EMLc)
<b>Formulations</b>	Oral > Liquid: 125 mg per 5 mL (as trihydrate) powder for oral liquid (EMLc) ; 250 mg per 5 mL (as trihydrate) powder for oral liquid (EMLc) Parenteral > General injections > unspecified: 250 mg in vial (as sodium) powder for injection ; 500 mg in vial (as sodium) powder for injection; 1 g in vial (as sodium) powder for injection Oral > Solid > dispersible tablet: 250 mg (scored) (as trihydrate) (EMLc) ; 500 mg (scored) (as trihydrate) (EMLc) Oral > Solid > dosage form: 250 mg (as trihydrate) ; 500 mg (as trihydrate)
<b>Historique des statuts LME</b>	Ajouté pour la première fois en 2017 (TRS 1006) Modifié en 2023 (TRS 1049)
<b>Sexe</b>	Tous
<b>Âge</b>	Aussi recommandé pour les enfants
<b>Équivalence thérapeutique</b>	La recommandation concerne ce médicament spécifique
<b>Renseignements sur le brevet</b>	Patents have expired in most jurisdictions Lire la suite sur les brevets.
<b>Wikipédia</b>	<a href="#">Amoxicillin</a>
<b>DrugBank</b>	<a href="#">Amoxicillin</a>

### Recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended: - the addition of new formulations of amoxicillin (dispersible, scored tablet, 250 mg and 500 mg) to the EMLc. - the addition of new formulations of cefalexin (dispersible tablet 125 mg and 250 mg) to the EMLc. - the addition of a new strength formulation of clarithromycin (solid oral dosage form 250 mg) to the EMLc. - the deletion of clarithromycin 500 mg solid oral dosage form from the EMLc.

#### Recommandations de la LME : Acute pharyngitis

##### Premier choix

##### Second choix

**WATCHFUL WAITING, SYMPTOM RELIEF AND NO ANTIBIOTIC TREATMENT SHOULD BE CONSIDERED AS THE FIRST-LINE TREATMENT OPTION.**

phenoxyethylpenicillin

cefalexin

amoxicillin

clarithromycin

