



EMLc

Codes ATC: J01DB01

<b>Indication</b>	Acute pharyngitis	Code ICD11: CA02
<b>INN</b>	Cefalexin	
<b>Type de médicament</b>	Chemical agent	
<b>Groupes d'antibiotiques</b>	<span style="border: 1px solid green; padding: 2px;">A ACCESS</span>	
<b>Type de liste</b>	Liste de base (EML) (EMLc)	
<b>Formulations</b>	Oral > Liquid: 125 mg per 5 mL (anhydrous) powder for oral liquid ; 250 mg per 5 mL (anhydrous) powder for oral liquid Oral > Solid: 250 mg (as monohydrate) ; 500 mg (as monohydrate) (EML) Oral > Solid > dispersible tablet: 125 mg (EMLc) ; 250 mg (EMLc)	
<b>Historique des statuts LME</b>	Ajouté pour la première fois en 2017 (TRS 1006) Modifié en 2021 (TRS 1035) Modifié en 2023 (TRS 1049)	
<b>Sexe</b>	Tous	
<b>Âge</b>	Aussi recommandé pour les enfants	
<b>Équivalence thérapeutique</b>	La recommandation concerne ce médicament spécifique	
<b>Renseignements sur le brevet</b>	Patents have expired in most jurisdictions Lire la suite <a href="#">sur les brevets</a> .	
<b>Wikipédia</b>	Cefalexin	
<b>DrugBank</b>	Cefalexin (Cephalexin)	

### Recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended: - the addition of new formulations of amoxicillin (dispersible, scored tablet, 250 mg and 500 mg) to the EMLc. - the addition of new formulations of cefalexin (dispersible tablet 125 mg and 250 mg) to the EMLc. - the addition of a new strength formulation of clarithromycin (solid oral dosage form 250 mg) to the EMLc. - the deletion of clarithromycin 500 mg solid oral dosage form from the EMLc.

### Recommandations de la LME : Acute pharyngitis

#### Premier choix

#### Second choix

**WATCHFUL WAITING, SYMPTOM RELIEF AND NO ANTIBIOTIC TREATMENT SHOULD BE CONSIDERED AS THE FIRST-LINE TREATMENT OPTION.**

phenoxycephalothin

cefalexin

amoxicillin

clarithromycin

