

EMLc

ATC codes: **N05CD08**

Indication	Status epilepticus ICD11 code: 8A66
INN	Midazolam
Medicine type	Chemical agent
List type	Core (EML) (EMLc)
Additional notes	Midazolam injection may be used for buccal administration when solution for oromucosal administration is not available.
Formulations	Parenteral > General injections > unspecified: 1 mg per mL in 5 mL vial (for buccal administration) ; 5 mg per mL in 1 mL vial (for buccal administration) ; 5 mg per mL in 3 mL vial (for buccal administration) Local > Buccal > Solution: 5 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration) ; 5 mg per mL in 1 mL pre-filled syringe (for oromucosal administration) ; 5 mg per mL in 1.5 mL pre-filled syringe (for oromucosal administration) ; 10 mg per mL in 0.25 mL pre-filled syringe (for oromucosal administration) ; 10 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration) ; 10 mg per mL in 0.75 mL pre-filled syringe ; 10 mg per mL in 1 mL pre-filled syringe (for oromucosal administration)
EML status history	First added in 2015 (TRS 994) Changed in 2023 (TRS 1049)
Sex	All
Age	Also recommended for children
Therapeutic alternatives	The recommendation is for this specific medicine
Patent information	Patents have expired in most jurisdictions Read more about patents . 
Wikipedia	Midazolam 
DrugBank	Midazolam 

Summary of evidence and Expert Committee recommendations

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended amendments to the listing for midazolam for status epilepticus on the EML and EMLc to better describe the available dosage forms.

