

		EMLc	Codes ATC: N05CD08
Indication	Status epilepticus	Code ICD11: 8A66	
INN	Midazolam		
Type de médicament	Chemical agent		
Type de liste	Liste de base (EML) (EMLc)		
Additional notes	Midazolam injection may be used for buccal administration when solution for oromucosal administration is not available.		
Formulations	Parenteral > General injections > unspecified: 1 mg per mL in 5 mL vial (for buccal administration) ; 5 mg per mL in 1 mL vial (for buccal administration) ; 5 mg per mL in 3 mL vial (for buccal administration) Local > Buccal > Solution: 5 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration) ; 5 mg per mL in 1 mL pre-filled syringe (for oromucosal administration) ; 5 mg per mL in 1.5 mL pre-filled syringe (for oromucosal administration) ; 5 mg per mL in 2 mL pre-filled syringe (for oromucosal administration) ; 10 mg per mL in 0.25 mL pre-filled syringe (for oromucosal administration) ; 10 mg per mL in 0.5 mL pre-filled syringe (for oromucosal administration) ; 10 mg per mL in 0.75 mL pre-filled syringe ; 10 mg per mL in 1 mL pre-filled syringe (for oromucosal administration)		
Historique des statuts LME	Ajouté pour la première fois en 2015 (TRS 994) Modifié en 2023 (TRS 1049)		
Sexe	Tous		
Âge	Aussi recommandé pour les enfants		
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique		
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets. 		
Wikipédia	Midazolam 		
DrugBank	Midazolam 		

Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended amendments to the listing for midazolam for status epilepticus on the EML and EMLc to better describe the available dosage forms.

