

[Ceftriaxone](#)

Essential medicine status

Section:

[6. Anti-infective medicines 6.2. Antibacterials 6.2.2. Watch group antibiotics](#)

ATC codes: [J01DD04](#)

EMLc

Indication

Infectious gastroenteritis or colitis without specification of infectious agent ICD11 code: [1A70.Z](#)

INN

Ceftriaxone

Medicine type

Chemical agent

Antibiotic groups

[WATCH](#)

List type

Core

Additional notes

Do not administer with calcium and avoid in infants with hyperbilirubinaemia.

Formulations

Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as sodium salt) ; 500 mg in vial powder for injection (as sodium salt) ; 1 g in vial powder for injection (as sodium salt) ; 2 g in vial powder for injection (as sodium salt) (EML)

EML status history

First added in 2017 ([TRS 1006](#))

Changed in 2023 ([TRS 1049](#))

Sex

All

Age

Also recommended for children

Age restriction

> 41 weeks corrected gestational age

Therapeutic alternatives

The recommendation is for this specific medicine

Patent information

Patents have expired in most jurisdictions

Read more [about patents](#).

Wikipedia

[Ceftriaxone](#)

DrugBank

[Ceftriaxone](#)

Expert Committee recommendation

In consideration of the review of the age appropriateness of formulations of medicines on the EMLc, and the comparison report of the EML versus EMLc, the Expert Committee recommended changes to the EMLc for addition of new, age-appropriate formulations and strengths of existing essential medicines, deletion of unavailable or age-inappropriate formulations and strengths, and other listing modifications as proposed in the application. The Committee also endorsed the proposals for further review of the public health relevance and evidence for specific medicines for use in children for potential future consideration for inclusion on the EMLc. The Committee noted and welcomed the ongoing review being coordinated by the Secretariat for the remaining sections of the EMLc for consideration by the 2025 Expert Committee. As a result of the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended: - replacing the oral liquid formulation of azithromycin (200 mg/5 mL) with powder for oral liquid 200 mg/5 mL on the EML and EMLc. - the addition of a new strength formulation of ceftriaxone (powder for injection: 500 mg (as sodium) in vial) to the EML and EMLc. - the addition of a new strength formulation of ciprofloxacin (solid oral dosage form 100 mg (as hydrochloride)) to the EMLc. - the addition of a new strength formulation of sulfamethoxazole + trimethoprim (dispersible tablet 100 mg + 20 mg) on the EMLc.

EML recommendations: Infectious gastroenteritis or colitis without specification of infectious agent

First choice

Second choice

First choice

[ciprofloxacin](#)

Second choice

[cefixime](#)

[azithromycin](#)

[sulfamethoxazole + trimethoprim](#)

[ceftriaxone](#)