

## [Amoxicillin + clavulanic acid](#)

Statut de médicament essentiel

Section:

[6. Anti-infective medicines 6.2. Antibacterials 6.2.1. Access group antibiotics](#)

Codes ATC: [J01CR02](#)

EMLc

Indication

Other specified pneumonia Code ICD11: [CA40.Y](#)

INN

Amoxicillin + clavulanic acid

Type de médicament

Chemical agent

Groupes d'antibiotiques

[ACCESS](#)

Type de liste

Liste de base

Formulations

**Parenteral > General injections > IV:** 500 mg (as sodium salt) + 100 mg (as potassium salt) powder for injection ; 1000 mg (as sodium salt) + 200 mg (as potassium salt) powder for injection

**Oral > Liquid:** 125 mg (as trihydrate) + 31.25 mg (as potassium salt) / 5 mL powder for oral liquid (EMLc) ; 250 mg (as trihydrate) + 62.5 mg (as potassium salt) / 5 mL powder for oral liquid (EMLc)

**Oral > Solid > dispersible tablet:** 200 mg (as trihydrate) + 28.5 mg (as potassium salt) (EMLc) ; 250 mg (as trihydrate) + 62.5 mg (as potassium salt) (EMLc)

**Oral > Solid > tablet:** 500 mg (as trihydrate) + 125 mg (as potassium salt) ; 875 mg (as trihydrate) + 125 mg (as potassium salt) (EML)

Historique des statuts LME

Ajouté pour la première fois en 2017 ([TRS 1006](#))

Modifié en 2023 ([TRS 1049](#))

Sexe

Tous

Âge

Aussi recommandé pour les enfants

Équivalence thérapeutique

La recommandation concerne ce médicament spécifique

Renseignements sur le brevet

Patents have expired in most jurisdictions

Lire la suite [sur les brevets.](#)

Wikipédia

[Amoxicillin + clavulanic acid](#)

DrugBank

[Amoxicillin](#),

[Clavulanic acid \(Clavulanate\)](#)

Recommandation du comité d'experts



The Expert Committee recommended the inclusion of a new strength, child-friendly dispersible tablet formulation of amoxicillin + clavulanic acid (200 mg + 28.5 mg) as an Access group antibiotic on the core list of the EMLc for treatment of bacterial infections in children - specifically those infections for which amoxicillin + clavulanic acid is already recommended on the EMLc. The Committee noted that the 7:1 ratio of amoxicillin to clavulanic acid is associated with similar efficacy to the 4:1 ratio but has a reduced frequency of gastrointestinal adverse effects. The Committee endorsed the importance of age-appropriate formulations to better meet the dosing needs of children. Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended: - the addition of a new formulation of amoxicillin + clavulanic acid (dispersible tablet 250 mg + 62.5 mg) to the EMLc. - the addition of new strength formulations of cefotaxime (powder for injection: 500 mg; 1 g; 2 g (as sodium) in vial) to the EML and EMLc. - the addition of a new strength formulation of ceftriaxone (powder for injection: 500 mg (as sodium) in vial) to the EML and EMLc.

Recommandations de la LME : Other specified pneumonia

Premier choix

Second choix

Hospital-acquired pneumonia

Premier choix

[piperacillin + tazobactam](#)

[amoxicillin + clavulanic acid](#)

[cefotaxime](#)

[ceftriaxone](#)

Second choix