## Ethambutol + isoniazid + pyrazinamide + rifampicin



Essential medicine status 🗸

Section: 6. Anti-infective medicines > 6.2. Antibacterials > 6.2.5. Antituberculosis medicines

	ATC codes: JC	)4AM0
Indication	Tuberculosis ICD11 code: 1B4Z	
INN	Ethambutol + isoniazid + pyrazinamide + rifampicin	
Medicine type	Chemical agent	
List type	Core	
Additional notes	WHO recommends and endorses the use of fixed-dose combinations and the development appropriate new fixed-dose combinations, including modified dosage forms, non-refriger products and paediatric dosage forms of assured pharmaceutical quality.	nt of ated
Formulations	Oral > Solid: 275 mg + 75 mg + 400 mg + 150 mg tablet	
EML status history	First added in 1999 (TRS 895) Changed in 2007 (TRS 950)	
Sex	All	
Age	Adolescents and adults	
Therapeutic alternatives	The recommendation is for this specific medicine	
Patent information	Patents have expired in most jurisdictions Read more about patents.	
Wikipedia	Ethambutol + isoniazid + pyrazinamide + rifampicin	
DrugBank	Ethambutol , soniazid , soniazid , Pyrazinamide , Rifampicin .	

## Summary of evidence and Expert Committee recommendations

The EMLc Subcommittee did not endorse the inclusion of this fixed-dose combination for the treatment of tuberculosis on the EMLc. The various fixed-dose combinations were considered as a group, as there is no clinical evidence for any of these combinations in children. However, it is difficult to determine the necessary appropriate combinations and what the strengths of the components should be in FDCs for use in children without examining further data (including pharmacokinetic data, stratified by weight and age). The Subcommittee therefore decided to endorse the lower strength rifampicin + isoniazid combinations (60 mg + 30 mg and 60 mg + 60 mg) and rifampicin + isoniazid + pyrazinamide (60 mg + 30 mg + 150 mg) as probably useful for many children, but requested an urgent review of all clinical evidence to support these and other potential combinations.

