

Section: 6. Anti-infective medicines > 6.2. Antibacterials > 6.2.2. Watch group antibiotics

	ATC codes: J01DD0				
Indication	Inflammatory and other diseases of prostate ICD11 code: GA91.Z				
INN	Cefotaxime				
Medicine type	Chemical agent				
Antibiotic groups	w WATCH				
List type	Core				
Additional notes	3rd generation cephalosporin of choice for use in hospitalized neonates.				
Formulations	Parenteral > General injections > unspecified: 250 mg in vial powder for injection (as s salt); 500 mg in vial powder for injection (as sodium salt); 1 g in vial powder for injecti sodium salt); 2 g in vial powder for injection (as sodium salt)				
EML status history	First added in 2017 (TRS 1006) Changed in 2023 (TRS 1049)				
Sex	Male				
Age	Adolescents and adults				
Therapeutic alternatives	The recommendation is for this specific medicine				
Patent information	Patents have expired in most jurisdictions Read more about patents.				
Wikipedia	Cefotaxime 🗹				
DrugBank	Cefotaxime 🗹				

Expert Committee recommendation

In consideration of the review of the age appropriateness of formulations of medicines on the EMLc, and the comparison report of the EML versus EMLc, the Expert Committee recommended changes to the EMLc for addition of new, age-appropriate formulations and strengths of existing essential medicines, deletion of unavailable or age-inappropriate formulations and strengths, and other listing modifications as proposed in the application. The Committee also endorsed the proposals for further review of the public health relevance and evidence for specific medicines for use in children for potential future consideration for inclusion on the EMLc. The Committee noted and welcomed the ongoing review being coordinated by the Secretariat for the remaining sections of the EMLc for consideration by the 2025 Expert Committee. As a result of the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended: - the addition of a new strength formulation of amikacin (injection: 500 mg/mL (as sulfate) in 2 mL vial) to the EMLc. - the addition of new strength formulation of ceftriaxone (powder for injection: 500 mg (as sodium) in vial) to the EML and EMLc. - the addition of a new strength formulation of ciprofloxacin (solid oral dosage form 100 mg (as hydrochloride)) to the EMLc.

Ŧ	d	V	IL recommend	lations.	Inflammat	ory and	ot	her d	iseases of	nrost	ate
•	-	ч	LD I CCOMMICHO	iutions.	IIIIIIIIII	.ory uno		IICI U	ilocuoco ol	PLOS	ucc

First choice	Second choice

MILD TO MODERATE

ciprofloxacin	cefotaxime
	ceftriaxone

SEVERE

cefotaxime co-prescribed with <u>amikacin</u>

ceftriaxone

 $\hbox{co-prescribed with } \underline{\hbox{amikacin}}$

