

Codes ATC: J01DD04

Indication	Inflammatory and other diseases of prostate	Code ICD11: GA91.Z
INN	Ceftriaxone	
Type de médicament	Chemical agent	
Groupes d'antibiotiques	 WATCH	
Type de liste	Liste de base	
Additional notes	Do not administer with calcium and avoid in infants with hyperbilirubinaemia.	
Formulations	Parenteral > General injections > unspecified: 2 g in vial powder for injection (as sodium salt) (EML) ; 500 mg in vial powder for injection (as sodium salt) ; 250 mg in vial powder for injection (as sodium salt) ; 1 g in vial powder for injection (as sodium salt)	
Historique des statuts LME	Ajouté pour la première fois en 2017 (TRS 1006) Modifié en 2023 (TRS 1049)	
Sexe	Mâle	
Âge	Adolescents et adultes	
Limite d'âge	> 41 weeks corrected gestational age	
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique	
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets .	
Wikipédia	Ceftriaxone	
DrugBank	Ceftriaxone	

Recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended: - the addition of a new strength formulation of amikacin (injection 50 mg/mL (as sulfate) in 2 mL vial) to the EMLc. - the addition of new strength formulations of cefotaxime (powder for injection: 500 mg; 1 g; 2 g (as sodium) in vial) to the EML and EMLc. - the addition of a new strength formulation of ceftriaxone (powder for injection: 500 mg (as sodium) in vial) to the EML and EMLc. - the addition of a new strength formulation of ciprofloxacin (solid oral dosage form 100 mg (as hydrochloride)) to the EMLc.

Recommandations de la LME : Inflammatory and other diseases of prostate

Premier choix

Second choix

MILD TO MODERATE

ciprofloxacin

ceftriaxone

cefotaxime

SEVERE

ceftriaxone

co-prescrite avec [amikacin](#)

cefotaxime

co-prescrite avec [amikacin](#)

