Chloramphenicol

Section: 6. Anti-infective medicines > 6.2. Antibacterials > 6.2.1. Access group antibiotics

EMLc | Codes ATC: J01BA01

Indication	Bacterial meningitis Code ICD11: 1D01.0Z
INN	Chloramphenicol
Type de médicament	Chemical agent
Groupes d'antibiotiques	A ACCESS
Type de liste	Liste de base
Additional notes	Oily suspension for injection formulation is only for the presumptive treatment of epidemic meningitis in children older than 2 years and in adults.
Formulations	**Parenteral > General injections > IV:** 1 g in vial powder for injection (sodium succinate) **Parenteral > General injections > IM:** 500 mg in 2 mL ampoule oily suspension for injection
Historique des statuts LME	Ajouté pour la première fois en 2017 (TRS 1006) Modifié en 2023 (TRS 1049)
Sexe	Tous
Âge	Aussi recommandé pour les enfants
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets.
Wikipédia	Chloramphenicol
DrugBank	Chloramphenicol

Recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended: - the addition of new strength formulations of cefotaxime (powder for injection: 500 mg; 1 g; 2 g (as sodium) in vial) to the EML and EMLc. - the addition of a new strength formulation of ceftriaxone (powder for injection: 500 mg (as sodium) in vial) to the EML and EMLc. - the addition of new formulations of amoxicillin (dispersible, scored tablet, 250 mg and 500 mg) to the EMLc. - the deletion of oral formulations of chloramphenicol (capsule 250 mg and oral liquid 150 mg/5 mL (as palmitate)) from the EML and EMLc.

Recommandations de la LME : Bacterial meningitis

Premier choix | **Second choix**

cefotaxime

ceftriaxone

cefotaxime

ceftriaxone

ampicillin

benzylpenicillin

amoxicillin

chloramphenicol

amoxicillin

ampicillin

benzylpenicillin

chloramphenicol

