

[Metronidazole](#)

Statut de médicament essentiel

Section:

[6. Anti-infective medicines 6.2. Antibacterials 6.2.1. Access group antibiotics](#)

Codes ATC: [A01AB17](#)

EMLc

Indication

Intestinal infections due to Clostridioides difficile Code ICD11: [1A04](#)

INN

Metronidazole

Type de médicament

Chemical agent

Groupes d'antibiotiques

[ACCESS](#)

Type de liste

Liste de base

Formulations

Oral > Liquid: 200 mg per 5 mL (as benzoate)

Local > Rectal > Suppository: 500 mg (EML) ; 1 g (EML)

Parenteral > General injections > unspecified: 500 mg per 100 mL in vial

Oral > Solid > tablet: 200 mg ; 250 mg ; 400 mg ; 500 mg

Historique des statuts LME

Ajouté pour la première fois en 2017 ([TRS 1006](#))

Modifié en 2023 ([TRS 1049](#))

Sexe

Tous

Âge

Aussi recommandé pour les enfants

Équivalence thérapeutique

La recommandation concerne ce médicament spécifique

Renseignements sur le brevet

Patents have expired in most jurisdictions

Lire la suite [sur les brevets.](#)

Wikipédia

[Metronidazole](#)

DrugBank

[Metronidazole](#)

Recommandation du comité d'experts



Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended a note be added to the listing of oral vancomycin on the EML and EMLc to indicate that the powder for injection dosage form may also be used for oral administration. Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended that the range of strengths of metronidazole tablets be replaced with reference to specific strengths on the EML and EMLc.

Recommandations de la LME : Intestinal infections due to Clostridioides difficile

Premier choix

Second choix

Premier choix

[metronidazole](#)

Second choix

[vancomycin](#)