




Section: [6. Anti-infective medicines](#) > [6.5. Antiprotozoal medicines](#) > [6.5.3. Antimalarial medicines](#) > [6.5.3.1. Antimalarial medicines](#) > For curative treatment

		EMLc	ATC codes: P01BA06
Indication	Malaria due to Plasmodium falciparum	ICD11 code: 1F40	
INN	Amodiaquine		
Medicine type	Chemical agent		
List type	Core (EML) (EMLc)		
Additional notes	To be used in combination with artesunate 50 mg		
Formulations	Oral > Solid: 153 mg (as hydrochloride) ; 200 mg (as hydrochloride)		
EML status history	First added in 1977 (TRS 615) Removed in 1979 (TRS 641) Added in 1982 (TRS 685) Changed in 1984 (TRS 722) Removed in 1987 (TRS 770) Added in 2003 (TRS 920) Changed in 2007 (TRS 946) Changed in 2007 (TRS 950) Changed in 2011 (TRS 965)		
Sex	All		
Age	Also recommended for children		
Therapeutic alternatives	The recommendation is for this specific medicine		
Patent information	Patents have expired in most jurisdictions Read more about patents . 		
Wikipedia	Amodiaquine 		
DrugBank	Amodiaquine 		

Summary of evidence and Expert Committee recommendations

Note specifying that amodiaquine alone can be used for the treatment of *P. vivax*, *P. ovale*, and *P. malariae* was deleted.
Amodiaquine should be used in combination with artesunate 50 mg.

