




| | | EMLc | ATC codes: P01BA06 |
|--------------------------|---|----------------------------------|------------------------------------|
| Indication | Malaria due to Plasmodium falciparum | ICD11 code: 1F40 | |
| INN | Amodiaquine | | |
| Medicine type | Chemical agent | | |
| List type | Core (EML) (EMLc) | | |
| Additional notes | To be used in combination with artesunate 50 mg | | |
| Formulations | Oral > Solid: 153 mg (as hydrochloride) ; 200 mg (as hydrochloride) | | |
| EML status history | First added in 1977 (TRS 615) Removed in 1979 (TRS 641) Added in 1982 (TRS 685) Changed in 1984 (TRS 722) Removed in 1987 (TRS 770) Added in 2003 (TRS 920) Changed in 2007 (TRS 946) Changed in 2007 (TRS 950) Changed in 2011 (TRS 965) | | |
| Sex | All | | |
| Age | Also recommended for children | | |
| Therapeutic alternatives | The recommendation is for this specific medicine | | |
| Patent information | Patents have expired in most jurisdictions Read more about patents .  | | |
| Wikipedia | Amodiaquine  | | |
| DrugBank | Amodiaquine  | | |

Summary of evidence and Expert Committee recommendations

Note specifying that amodiaquine alone can be used for the treatment of *P. vivax*, *P. ovale*, and *P. malariae* was deleted.
Amodiaquine should be used in combination with artesunate 50 mg.

