Amodiaquine

NOT RECOMMENDED AS AN ESSENTIAL MEDICINE

Section: 6. Anti-infective medicines > 6.5. Antiprotozoal medicines > 6.5.3. Antimalarial medicines > 6.5.3.1. Antimalarial medicines > Medicines for curative treatment

EMLc Codes ATC: P01BA06

Indication	Malaria due to Plasmodium falciparum Code ICD11: 1F40
INN	Amodiaquine
Type de médicament	Chemical agent
Type de liste	Liste de base
Additional notes	To be used in combination with artesunate 50 mg
Formulations	**Oral > Solid:** 153 mg (as hydrochloride) ; 200 mg (as hydrochloride)
Historique des statuts LME	Ajouté pour la première fois en 1977 (TRS 615) Retiré en 1979 (TRS 641) Ajouté en 1982 (TRS 685) Modifié en 1984 (TRS 722) Retiré en 1987 (TRS 770) Ajouté en 2003 (TRS 920) Modifié en 2007 (TRS 946) Modifié en 2007 (TRS 950) Modifié en 2011 (TRS 965) Retiré en 2025 (TRS 1064)
Sexe	Tous
Âge	Aussi recommandé pour les enfants
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets.
Wikipédia	Amodiaquine
DrugBank	Amodiaquine

Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations of essential medicines for children, the Expert Committee recommended the removal from the EML and EMLc of single-agent amodiaquine tablets. These formulations were not found registered at stringent regulatory authorities. Fixed-dose combination formulations of amodiaquine and artesunate are preferred and are already included on the Model Lists.

