Artemether

Section: 6. Anti-infective medicines > 6.5. Antiprotozoal medicines > 6.5.3. Antimalarial medicines > 6.5.3.1. Antimalarial medicines > Medicines for curative treatment

EMLc | Codes ATC: P01BE02

Indication	Malaria due to Plasmodium falciparum Code ICD11: 1F40
INN	Artemether
Type de médicament	Chemical agent
Type de liste	Liste de base
Additional notes	For use in the management of severe malaria
Formulations	**Parenteral > General injections > IM:** 20 mg per mL in 1 mL ampoule (oily injection) ; 40 mg per mL in 1 mL ampoule (oily injection)
Historique des statuts LME	Ajouté pour la première fois en 1997 (TRS 882) Modifié en 2002 (TRS 914) Modifié en 2007 (TRS 946) Modifié en 2007 (TRS 950) Modifié en 2025 (TRS 1064)
Sexe	Tous
Âge	Aussi recommandé pour les enfants
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets.
Wikipédia	Artemether
DrugBank	Artemether

Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations of essential medicines for children, the Expert Committee recommended replacing the 80 mg/mL strength of artemether oily injection with 20 mg/mL and 40 mg/mL strengths. The 80 mg/mL strength has limited utility for children under 12 years of age.