

## [Mefloquine](#)

Not recommended as an essential medicine

Section:

[6. Anti-infective medicines 6.5. Antiprotozoal medicines 6.5.3. Antimalarial medicines 6.5.3.1. Antimalarial medicines > Medicines for curative treatment](#)

ATC codes: [P01BC02](#)

EMLc

Indication

Malaria due to Plasmodium falciparum ICD11 code: [1F40](#)

INN

Mefloquine

Medicine type

Chemical agent

List type

Core

Additional notes

To be used in combination with artesunate 50 mg

Formulations

**Oral > Solid:** 250 mg tablet (hydrochloride)

EML status history

First added in 1987 ([TRS 770](#))

Changed in 1993 ([TRS 850](#))

Changed in 2003 ([TRS 920](#))

Changed in 2007 ([TRS 946](#))

Changed in 2007 ([TRS 950](#))

Removed in 2025 ([TRS 1064](#))

Sex

All

Age

Also recommended for children

Therapeutic alternatives

The recommendation is for this specific medicine

Patent information

Patents have expired in most jurisdictions

Read more [about patents](#).

Wikipedia

[Mefloquine](#)

DrugBank

[Mefloquine](#)

Summary of evidence and Expert Committee recommendations

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended: - to delete the listing of single agent mefloquine from the EML and EMLc, noting that fixed dose combinations of artesunate + mefloquine are available and preferred for the curative treatment of malaria.