




Section: [6. Anti-infective medicines](#) > [6.5. Antiprotozoal medicines](#) > [6.5.3. Antimalarial medicines](#) > [6.5.3.1. Antimalarial medicines](#) > For curative treatment

		EMLc	ATC codes: <a href="#">P01BC01</a>
Indication	Malaria due to Plasmodium falciparum	ICD11 code: <a href="#">1F40</a>	
Medicine type	Chemical agent		
List type	Core (EML) (EMLc)		
Additional notes	For use only in the management of severe malaria, and should be used in combination with doxycycline.		
Formulations	Parenteral > General injections > IV: 300 mg per mL in 2 mL ampoule (quinine hydrochloride) Oral > Solid: 300 mg (quinine sulfate or quinine bisulfate)		
EML status history	First added in 1977 ( <a href="#">TRS 615</a> ) Changed in 1979 ( <a href="#">TRS 641</a> ) Changed in 1982 ( <a href="#">TRS 685</a> ) Changed in 1987 ( <a href="#">TRS 770</a> ) Changed in 2003 ( <a href="#">TRS 920</a> ) Changed in 2007 ( <a href="#">TRS 946</a> ) Changed in 2007 ( <a href="#">TRS 950</a> )		
Sex	All		
Age	Also recommended for children		
Therapeutic alternatives	The recommendation is for this specific medicine		
Patent information	Patents have expired in most jurisdictions Read more <a href="#">about patents</a> . 		
Wikipedia	<a href="#">Quinine</a> 		
DrugBank	<a href="#">Quinine</a> 		

### Summary of evidence and Expert Committee recommendations

Quinine was added to the core list of the EMLc for the curative treatment of malaria.

