Suramin sodium

Section: 6. Anti-infective medicines > 6.5. Antiprotozoal medicines > 6.5.5. Antitrypanosomal medicines > 6.5.5.1. African trypanosomiasis > 6.5.5.1.1. Medicines for the treatment of 1st stage African trypanosomiasis

EMLc | Codes ATC: P01CX02

Indication	Rhodesiense trypanosomiasis Code ICD11: 1F51.1
INN	Suramin sodium
Type de médicament	Chemical agent
Type de liste	Liste de base
Additional notes	To be used for the treatment of the initial phase of Trypanosoma brucei rhodesiense infection.
Formulations	**Parenteral > General injections > IV:** 1 g in vial
Historique des statuts LME	Ajouté pour la première fois en 1977 (TRS 615) Modifié en 1979 (TRS 641) Modifié en 2007 (TRS 946) Modifié en 2007 (TRS 950)
Sexe	Tous
Âge	Aussi recommandé pour les enfants
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets.
Wikipédia	Suramin sodium
DrugBank	Suramin sodium (Suramin)

Résumé des preuves et recommandation du comité d'experts

The EMLc Subcommittee recommended that suramin sodium be endorsed for the EMLc for treatment of first-stage of Trypanosoma brucei rhodesiense infections. The Subcommittee noted that African trypanosomiasis is a major public health problem in affected countries. Medicines available for treating this infection are limited. Hence despite the toxicity profile, these medicines are still used. There is a paucity of documented information specific to children and recommendations for adults are usually followed.