




Section: [6. Anti-infective medicines](#) > [6.5. Antiprotozoal medicines](#) > [6.5.5. Antitrypanosomal medicines](#) > [6.5.5.1. African trypanosomiasis](#) > [6.5.5.1.2. Medicines for the treatment of 2nd stage African trypanosomiasis](#)

EMLc

Codes ATC: **P01CX03**

Indication	African trypanosomiasis Code ICD11: 1F51
INN	Eflornithine
Type de médicament	Chemical agent
Type de liste	Liste de base (EML) (EMLc)
Additional notes	To be used for the treatment of <i>Trypanosoma brucei gambiense</i> infection.
Formulations	Parenteral > General injections > IV: 200 mg per mL in 50 mL bottle (hydrochloride)
Historique des statuts LME	Ajouté pour la première fois en 1991 (TRS 825) Modifié en 2007 (TRS 946) Modifié en 2007 (TRS 950) Modifié en 2023 (TRS 1049)
Sexe	Tous
Âge	Aussi recommandé pour les enfants
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets. 
Wikipédia	Eflornithine 
DrugBank	Eflornithine 

Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended the listing for eflornithine on the EML and EMLc be amended to change the bottle volume in which eflornithine is supplied from 100 mL to 50 mL.

