

## [Ceftazidime + avibactam](#)

Statut de médicament essentiel

Section:

[6. Anti-infective medicines 6.2. Antibacterials 6.2.3. Reserve group antibiotics](#)

Codes ATC: [J01DD52](#)

EMLc

Indication

Carbapenem-resistant *Pseudomonas aeruginosa* Code ICD11: [MG50.80](#)

INN

Ceftazidime + avibactam

Type de médicament

Chemical agent

Groupes d'antibiotiques

[RESERVE](#)

Type de liste

Liste complémentaire

Formulations

**Parenteral > General injections > IV:** 2 g in vial + 0.5 g in vial powder for injection

Historique des statuts LME

Ajouté pour la première fois en 2019 ([TRS 1021](#))

Sexe

Tous

Âge

Aussi recommandé pour les enfants

Équivalence thérapeutique

La recommandation concerne ce médicament spécifique

Renseignements sur le brevet

Main patent is active in several jurisdictions. For more information on specific patents and license status for developing countries visit [www.MedsPal.org](http://www.MedsPal.org)

Lire la suite [sur les brevets.](#)

Wikipédia

[Ceftazidime + avibactam](#)

DrugBank

[Ceftazidime](#),

[Avibactam](#)

Recommandation du comité d'experts

The Expert Committee recommended the inclusion of ceftazidime + avibactam on the complementary list of the EML and EMLc for the treatment of infections caused by carbapenem-resistant organisms which are pathogens classified as “critical priority” in the WHO Priority Pathogen List. The Committee agreed with the EML Antibiotic Working Group’s recommendation that this antibiotic should be classified in the AWaRe Reserve group. The Committee recommended further collaboration between relevant stakeholders to design and implement strategic public health orientated studies that will help to inform the choice of optimal single or combination treatment of both novel and older antibiotics for adults and children in different settings, with the goal of improving clinical outcomes, minimizing toxicity and reducing selection of resistance.

Contexte

This combination antibiotic had not previously been considered for inclusion on the EML. Ceftazidime is third generation cephalosporin listed on the EML complementary list and classified within the Watch group. Avibactam is a non-beta-lactam beta-lactamase inhibitor active against certain types of carbapenemases (e.g. KPC and OXA-48 but not active against metallo-beta-lactamases).

Pertinence pour la santé publique

Antibiotic-resistant bacteria are a significant threat to public health, both in high-income as well as low- and middle-income countries (LMIC) (1-3). A recent study estimated that infections with antibiotic-resistant bacteria were responsible for approximately 33 000 attributable deaths in Europe in 2015 (1). Fewer data are available for LMIC, but a retrospective study in ten hospitals in India found that resistant pathogens were associated with 2-3 times higher mortality than infections with susceptible strains after adjusting for several confounders (2). Over the last decade there has been increasing spread of multidrug-resistant Gram-negative pathogens such as carbapenemase producing Enterobacteriaceae (4). The Global Antimicrobial Resistance Surveillance System (GLASS) Report published in 2018 found high levels of carbapenem resistance in Enterobacteriaceae and non-fermenters in many of the LMICs providing data for the report (2). The 2015 WHO Global Action Plan on Antimicrobial Resistance calls for the development of new antimicrobial medicines (3). To provide a framework for this endeavour, in 2017 WHO published a priority list of antibiotic-resistant bacteria – the WHO Priority Pathogens List (5). “Priority 1: critical” category includes four types of pathogens, all of which are Gram-negative: carbapenem resistant *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and Enterobacteriaceae; and third-generation cephalosporin resistant Enterobacteriaceae (6).

Bénéfices

Several RCTs have been conducted comparing ceftazidime + avibactam to carbapenems or best available therapy for cIAIs and cUTIs (7-10). Of note, all but one of the RCTs (7) included patients based on clinical syndromes and not based on the presence of infections confirmed to be caused by multidrug-resistant organisms. In that “descriptive” trial of patients with cUTI (plus some patients with cIAI) caused by ceftazidime-resistant Gram negatives, ceftazidime + avibactam treatment resulted in similar clinical response compared to best available therapy. So far, few data on the “real life” clinical use of ceftazidime + avibactam have been published. A retrospective single centre study at the

University of Pittsburgh Medical Centre in Pittsburgh, USA examined outcomes of 109 patients with bacteraemia caused by carbapenem-resistant *K. pneumoniae* bacteraemia (97% of which were KPC producers) over the time period from 2009 to 2017. The 30-day survival rate was 92% (12/13) in patients treated with ceftazidime + avibactam vs. 69% (66/96) for patients treated with other regimens, but this obviously has to be interpreted with caution given the many potential confounding factors (11). Published data about use of ceftazidime + avibactam in children is very scarce and limited to a phase I study and case reports (12-14). However, two phase 2 RCTs have been conducted in children with cUTIs and cIAI and are awaiting publication (ClinicalTrials.gov Identifier: NCT02475733 and NCT02497781). Of note ceftazidime + avibactam may have a role in combination with aztreonam to treat infections caused by Enterobacteriaceae producing metallo-beta-lactamases at least until the combination of aztreonam with avibactam becomes available (15, 16).

Torts



In the RCTs the incidence of adverse events in the groups treated with ceftazidime + avibactam was similar to the control groups (7-10). However, in a meta-analysis of eight RCTs including 4093 patients, serious adverse events (SAEs) were more common with ceftazidime + avibactam (RR 1.24, 95%CI 1.00-1.54, I<sup>2</sup> = 0%) but detailed data regarding the nature of these SAE were not available (17).

Preuves supplémentaires



N/A

Rapport coût/efficacité



United Kingdom: Basic NHS price: 10 vial pack £857.00 = £257.1 (about 340 USD) per day (standard dosing) Few data are available regarding the cost-effectiveness of ceftazidime-avibactam. A decision analytic model presented at ID week in October 2018 aimed to estimate the cost-effectiveness of treatment with ceftazidime + avibactam compared with colistin for a hypothetical cohort of patients with pneumonia and bacteraemia caused by carbapenemase-resistant Enterobacteriaceae over a 12 months period. The researchers assumed a 41% mortality with colistin treatment, a 23% (and hence very large) absolute reduction in mortality with ceftazidime-avibactam, daily costs of ceftazidime-avibactam of 1080 USD, a 42% incidence of nephrotoxicity with colistin treatment, a 56% probability of transfer to long-term care and a 1.8 fold improved odds of discharge home with ceftazidime-avibactam treatment (18). The authors estimated an incremental cost-effectiveness ratio for ceftazidime + avibactam compared with colistin of 110, 300 USD per quality adjusted life-year.

Directives de l'OMS



There are no available WHO guidelines for the treatment of infections due to multi-drug resistant organisms.

Disponibilité



Ceftazidime-avibactam has FDA & EMA approval for cUTI & cIAI (for cIAI in combination with metronidazole) (11). EMA lists "HAP and other infections due to Gram-negative bacteria with limited treatment options" as further indication.

Autres considérations



The Committee noted that there was very limited clinical trial evidence of the efficacy of recently approved antibiotics for infections caused by carbapenem-resistant bacteria, with activity against this type of infection based on studies with small sample sizes, methodological limitations and including heterogenous populations. The Committee was concerned that the current regulatory approval process for novel agents effective against "critical priority" pathogens (according to the WHO Priority Pathogens List (5)) does not adequately inform the urgent public health need for clear evidence-based guidance on the optimal management of these infections, which are associated with important morbidity and mortality.

Afficher les références Masquer les références

1. Cassini A, Hogberg LD, Plachouras D, Quattrocchi A, Hoxha A, Simonsen GS, et al. Attributable deaths and disability-adjusted life-years caused by infections with antibiotic-resistant bacteria in the EU and the European Economic Area in 2015: a population-level modelling analysis. *Lancet Infect Dis*. 2018.
2. Gandra S, Tseng KK, Arora A, Bhowmik B, Robinson ML, Panigrahi B, et al. The mortality burden of multidrug-resistant pathogens in India: a retrospective observational study. *Clin Infect Dis*. 2018.
3. World Health Organization. Global Action Plan on Antimicrobial Resistance. 2015.
4. van Duin D, Doi Y. The global epidemiology of carbapenemase-producing Enterobacteriaceae. *Virulence*. 2017;8(4):460-9.
5. Prioritization of pathogens to guide discovery, research and development of new antibiotics for drug-resistant bacterial infections, including tuberculosis. Geneva: World Health Organization; 2017(WHO/EMP/IAU/2017.12). (Licence: CC BY-NC-SA 3.0 IGO). Available from <https://apps.who.int/iris/handle/10665/311820>.
6. Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, et al. Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis. *Lancet Infect Dis*. 2018;18(3):318-27.
7. Carmeli Y, Armstrong J, Laud PJ, Newell P, Stone G, Wardman A, et al. Ceftazidime-avibactam or best available therapy in patients with ceftazidime-resistant Enterobacteriaceae and *Pseudomonas aeruginosa* complicated urinary tract infections or complicated intra-abdominal infections (REPRISE): a randomised, pathogen-directed, phase 3 study. *Lancet Infect Dis*. 2016;16(6):661-73.
8. Mazuski JE, Gasink LB, Armstrong J, Broadhurst H, Stone GG, Rank D, et al. Efficacy and Safety of Ceftazidime-Avibactam Plus Metronidazole Versus Meropenem in the Treatment of Complicated Intra-abdominal Infection: Results From a Randomized, Controlled, Double-Blind, Phase 3 Program. *Clin Infect Dis*. 2016;62(11):1380-9.
9. Qin X, Tran BG, Kim MJ, Wang L, Nguyen DA, Chen Q, et al. A randomised, double-blind, phase 3 study comparing the efficacy and safety of ceftazidime/avibactam plus metronidazole versus meropenem for complicated intra-abdominal infections in hospitalised adults in Asia. *Int J Antimicrob Agents*. 2017;49(5):579-88.
10. Wagenlehner FM, Sobel JD, Newell P, Armstrong J, Huang X, Stone GG, et al. Ceftazidime-avibactam Versus Doripenem for the Treatment of Complicated Urinary Tract Infections, Including Acute Pyelonephritis: RECAPTURE, a Phase 3 Randomized Trial Program. *Clin Infect Dis*. 2016;63(6):754-62.
11. Pogue JM, Bonomo RA, Kaye KS. Ceftazidime/avibactam, Meropenem/vaborbactam or both? Clinical and formulary considerations. *Clin Infect Dis*. 2018.
12. Bradley JS, Armstrong J, Arrieta A, Bishai R, Das

S, Delair S, et al. Phase I Study Assessing the Pharmacokinetic Profile, Safety, and Tolerability of a Single Dose of Ceftazidime-Avibactam in Hospitalized Pediatric Patients. *Antimicrob Agents Chemother.* 2016;60(10):6252-9. 13. Rodriguez BA, Girotto JE, Nicolau DP. Ceftazidime/Avibactam and Ceftolozane/Tazobactam: Novel Therapy for Multidrug Resistant Gram Negative Infections in Children. *Curr Pediatr Rev.* 2018;14(2):97-109. 14. Tamma PD, Fan Y, Bergman Y, Sick-Samuels AC, Hsu AJ, Timp W, et al. Successful Treatment of Persistent Burkholderia cepacia Complex Bacteremia with Ceftazidime-Avibactam. *Antimicrob Agents Chemother.* 2018;62(4). 15. Hobson CA, Bonacorsi S, Fahd M, Baruchel A, Cointe A, Poey N, et al. Successful treatment of a bacteremia due to NDM-1-producing Morganella morganii with Aztreonam and Ceftazidime-avibactam combination in a pediatric patient with hematologic malignancy. *Antimicrob Agents Chemother.* 2018. 16. Barlow G, Morice A. Successful treatment of resistant Burkholderia multivorans infection in a patient with cystic fibrosis using ceftazidime/avibactam plus aztreonam. *J Antimicrob Chemother.* 2018;73(8):2270-1. 17. Sternbach N, Leibovici Weissman Y, Avni T, Yahav D. Efficacy and safety of ceftazidime/avibactam: a systematic review and meta-analysis. *J Antimicrob Chemother.* 2018;73(8):2021-9. 18. Sfeir M, Satlin M, Calfee DP, Simon MS. Cost-Effectiveness of Ceftazidime-Avibactam Compared With Colistin for Treatment of Carbapenem-Resistant Enterobacteriaceae Bacteremia and Pneumonia. *Open Forum Infect Dis.* 2018;5: S539-S40.