




Section: 20. Muscle relaxants (peripherally-acting) and cholinesterase inhibitors

		EMLc	Codes ATC: N07AA01
Indication	Myasthenia gravis	Code ICD11: 8C70.Z	
INN	Neostigmine bromide		
Type de médicament	Chemical agent		
Type de liste	Liste de base (EML) (EMLc)		
Formulations	Parenteral > General injections > unspecified: 0.5 mg per mL in 1 mL ampoule (neostigmine metilsulfate) ; 2.5 mg per mL in 1 mL ampoule (neostigmine metilsulfate) Oral > Solid > tablet: 15 mg (neostigmine bromide) (EML)		
Historique des statuts LME	Ajouté pour la première fois en 1977 (TRS 615) Modifié en 1979 (TRS 641) Modifié en 1989 (TRS 796) Modifié en 2003 (TRS 920) Modifié en 2007 (TRS 950) Modifié en 2025 (TRS 1064)		
Sexe	Tous		
Âge	Aussi recommandé pour les enfants		
Équivalence thérapeutique	La recommandation concerne ce médicament spécifique		
Renseignements sur le brevet	Patents have expired in most jurisdictions Lire la suite sur les brevets. 		
Wikipédia	Neostigmine 		
DrugBank	Neostigmine 		

Résumé des preuves et recommandation du comité d'experts

Following the review of the age-appropriateness of formulations on the EMLc, the Expert Committee recommended: - the deletion of neostigmine 15 mg tablets from the EMLc

